

## RECENTLY DISCOVERED PULSARS AND UNIDENTIFIED EGRET SOURCES

DIEGO F. TORRES<sup>a</sup>, YOUSAF M. BUTT<sup>b</sup>, AND FERNANDO CAMILO<sup>c</sup>

<sup>a</sup>Physics Department, Princeton University, NJ 08544

<sup>b</sup>Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

<sup>c</sup>Columbia Astrophysics Laboratory, Columbia University, 550 West 120th Street, New York, NY 10027

*Accepted for publication in The Astrophysical Journal Letters*

### ABSTRACT

We present a correlative study between all unidentified EGRET sources at low Galactic latitudes and the newly discovered pulsars in the released portion of the Parkes multibeam radio survey. We note 14 positional coincidences: eight of these are “Vela-like” pulsars, with relatively small periods, small characteristic ages, and high spin-down luminosities. Three of these coincidences have been investigated by D’Amico et al. (2001) and Camilo et al. (2001). Among the others, we argue that PSR J1015–5719 may plausibly generate part of the high energy radiation observed from 3EG J1014–5705. Three additional interesting cases are: 3EG J1410–6147 and either of PSRs J1412–6145 or J1413–6141, if the pulsars are at the estimated distance of the coincident SNR G312.4–0.4; and 3EG J1639–4702/PSR J1637–4642. The remaining positional coincidences between the EGRET sources and the newly discovered pulsars are almost certainly spurious.

*Subject headings:* pulsars: general — pulsars: individual (PSR J1015–5719, PSR J1412–6145, PSR J1413–6141, PSR J1637–4642) — gamma rays: observations

### 1. INTRODUCTION

The Third EGRET Catalog lists 271 point sources (Hartman et al. 1999) of which about two thirds are not yet identified. Although most known young pulsars are not detected in the high-energy window ( $> 100$  MeV), there are so far six such confirmed  $\gamma$ -ray pulsars (Thompson 1999). They are relatively young objects, with hard high-energy spectral indices, and apparently showing a trend for spectral hardening with increasing characteristic age (Fierro et al. 1993). Young pulsars were proposed as potential counterparts for  $\gamma$ -ray sources as early as 1983 (D’Amico 1983; Helfand 1994), and their relationship with unidentified EGRET sources remains a matter of current debate (see, e.g., Zhang, Zhang, & Cheng 2000).

In order to unambiguously identify a pulsar as the origin of the  $\gamma$ -rays from an EGRET source,  $\gamma$ -ray pulsations must be detected at the pulsar period. However the  $\gamma$ -fluxes are generally very low, and nearly contemporaneous radio/X-ray and  $\gamma$ -ray observations are required to fold the few available photons with the correct ephemeris. Alas this is no longer possible owing to the demise of the *Compton Gamma-Ray Observatory*, and new candidate associations must for now be judged by comparison with the properties of the six known EGRET pulsars.

Supernova remnants (SNRs) interacting with nearby molecular clouds have also been proposed as counterparts to some  $\gamma$ -ray sources (e.g. Sturmer, Dermer, & Mattox 1996; Esposito et al. 1996; Combi et al. 1998a, b, 2001), and in such cases both a pulsar and an associated interacting SNR may partially contribute to generate a given EGRET source flux.

The Parkes multibeam pulsar survey is a large-scale survey of a narrow strip of the inner Galactic plane ( $|b| < 5^\circ$ ,  $260^\circ < l < 50^\circ$ ; Manchester et al. 2001). It has much greater sensitivity than any previous survey to young and distant pulsars along the Galactic plane, and it has resulted in the detection of many previously unknown young pulsars, potentially counterparts of EGRET sources. In this Letter we correlate the re-

leased portion of the Parkes pulsar survey<sup>1</sup>, now consisting of 368 pulsars, with all unidentified EGRET sources at low latitudes, in order to determine whether there are any new likely physical associations.

### 2. CORRELATIONS

Table 1 lists some important properties for the known EGRET  $\gamma$ -ray pulsars. Two of the parameters listed are highly uncertain: the distance  $d$ , and the *observed* efficiency for conversion of spin-down luminosity  $\dot{E}$  into  $\gamma$ -ray luminosity  $L_\gamma$ :  $\eta \equiv L_\gamma/\dot{E} = f4\pi d^2 F_\gamma/\dot{E}$ , where  $F_\gamma$  is the observed  $\gamma$ -ray flux and  $f$  is the  $\gamma$ -ray beaming fraction ( $0 < f \leq 1$ ). Because this fraction is essentially unknown (see, e.g., Yadigaroglu & Romani 1995), and in part because  $f \sim 1$  results in  $\eta \sim 100\%$  for at least one pulsar at its nominal distance, it is common practice, which we follow, to *assume*  $f \equiv 1/4\pi$ . The efficiencies observed span the range  $0.01\% \lesssim \eta \lesssim 3/19\%$ : the two upper limits for  $\eta$  stem from the different measurements of the distance to PSR B1055–52 (see Ögelman & Finley 1993; Combi et al. 1997; Romero 1998). We shall then consider  $0.01\% \lesssim \eta \lesssim 10\%$  as “reasonable” values for evaluating the plausibility of new associations, by analogy with presently known  $\gamma$ -ray pulsars, even though this inferred range relies on an assumed uniform value for  $f$ , and uncertain distances.

All positional coincidences between unidentified 3EG sources and Parkes pulsars are given in Table 2. We list for each 3EG source the variability index  $I$ : pulsars are steady  $\gamma$ -ray sources. For each pulsar we also present the “spin-down flux”  $\dot{E}/d^2$ , which is generally well-correlated with the detectability of  $\gamma$ -ray pulsars. A strict (linear) reading of this correlation would suggest  $L_\gamma \propto \dot{E}$ , while Thompson et al. (1999) find that  $L_\gamma \propto \dot{E}^{1/2}$ . However, we summarily dismiss some potential associations below owing to a low value of  $\dot{E}/d^2$  only in extreme cases where the difference between  $\dot{E}$ - and  $\dot{E}^{1/2}$ -scaling is irrelevant (and where even a hefty error in a more relevant and

<sup>1</sup> See <http://www.atnf.csiro.au/research/pulsar/pmsurv>.

uncertain parameter — distance — does not change matters significantly). In all these discarded cases, in addition, the efficiencies required would be  $\eta \gg 1000\%$ , making the potential associations utterly unphysical. Finally, in evaluating possible associations, we compare the photon indices of known pulsars (Table 1) with those of the EGRET sources (see, e.g., Merck et al. 1996; Zhang & Cheng 1998; Cheng & Zhang 1998).

We have also quantified the chance probability for obtaining these associations, adapting the numerical code described by Romero et al. (1999a, b): there is a 2% probability of having 8 chance coincidences between different unidentified 3EG sources and Parkes pulsars, as in Table 2.<sup>2</sup>

Camilo et al. (2001) proposed the possible physical association #1, with required efficiency of 0.5% (Table 2). This pulsar appears to be located just outside the SNR G284.3–1.8, which itself is interacting with an adjacent molecular cloud (Ruiz & May 1986). Another Parkes pulsar, PSR J1013–5934 (#2 in Table 2), is coincident with 3EG 1013–5915, but it is an old pulsar ( $\tau = 12$  Myr) with low  $\dot{E} = 2.5 \times 10^{32}$  ergs s<sup>-1</sup> and cannot be a significant  $\gamma$ -ray contributor. D’Amico et al. (2001) have studied cases #6 and 14 (efficiencies:  $\eta = 2\%$  and  $7\%$ , respectively). Both are plausible candidates to generate the respective EGRET source fluxes. PSR J1837–0559, is also coincident with 3EG J1837–0606 (#13), but its  $\dot{E}/d^2$  is 80 times smaller than for PSR J1837–0604, and it cannot contribute significantly to the  $\gamma$ -ray source. The pulsars in pairs #8 and 9 have far too low a spin-down luminosity at too large a distance (Table 2) to explain their coincident  $\gamma$ -ray sources. They are also both old, with  $\tau \sim 4$  Myr. Cases #10 and 11 are discussed elsewhere in connection with a proposal for a SNR shock origin of the bulk of the  $\gamma$ -rays resulting from 3EG J1714–3857 (Butt et al. 2001); neither of these pulsars is energetic enough to contribute significant amounts of high-energy flux.

### 3. DISCUSSION

We now discuss the remaining four EGRET sources positionally superposed with five newly discovered pulsars. We provide observational data on the apparent associations in Tables 3 and 4. Cases #3–5, 7 and 12 all contain Vela-like pulsars, with relatively short periods, low characteristic ages, and high spin-down luminosities ( $\dot{E} \gtrsim 10^{35}$  ergs s<sup>-1</sup>).

The pulsar in case #3 would require an efficiency  $\eta = 5\%$  at its nominal distance to explain the luminosity of the corresponding 3EG source, which has photon index 2.23 (see Table 4). This spectrum is softer than that of the Crab, although it is consistent with it within the uncertainties. It is also consistent with the index of 3EG 2227+6122, for which PSR J2229+6114 has been proposed as the likely source (Halpern et al. 2001). The  $\gamma$ -ray source in case #3 is not variable (Tompkins 1999; Torres et al. 2001c), as expected from direct pulsar or pulsar wind nebula/SNR shock emission. PSR J1015–5719 has  $\tau = 39$  kyr and  $\dot{E} = 8.2 \times 10^{35}$  ergs s<sup>-1</sup> (Table 3). While no cataloged SNR is superposed with the 3EG source (Torres et al. 2001b), this absence does not mean that one does not exist, and further sensitive searches may prove fruitful. Thus the connection between 3EG J1014–5705 and PSR J1015–5719 appears plausible and is worth additional study.

The pulsars in pairs #4 and 5 require unreasonably high efficiencies at their nominal distances to explain the  $\gamma$ -ray flux from the corresponding EGRET source ( $\eta \gtrsim 100\%$ , Table 4).

However, both pulsars are located in the direction of the Centaurus arm, and it is known that in such directions the electron density/distance model of Taylor & Cordes (1993) can be unreliable, sometimes overestimating the distances by factors of up to  $\sim 4$  (see discussion in Camilo et al. 2001). Both pulsars are located (at least in projection) well within the boundaries of the incomplete shell SNR G312.4–0.4 (Caswell & Barnes 1985), to which Yadigaroglu & Romani (1997) estimate a  $\Sigma - D$  distance of 1.9 kpc. At this distance the required efficiencies for the pulsars in cases #4 and 5 would be 12% and 3%, respectively, which would make them considerably more plausible sources of the observed high energy emission. Furthermore, 3EG J1410–6147 has a photon index comparable to that of the Crab, and is not variable (Tables 1, 2 and 4). Pairs #4 and 5 therefore appear intriguing. However it should be noted that  $\Sigma - D$  distances are notoriously unreliable (e.g., for this very SNR, Caswell & Barnes 1985 and Case & Bhattacharya 1999 infer values in substantial disagreement both with each other and with that determined by Yadigaroglu & Romani 1997). Ideally, further observations of SNR G312.4–0.4 may indicate whether it shows signs of interaction with PSRs J1412–6145 or J1413–6141, and possibly constrain their distances. Depending on the actual distances, the  $\gamma$ -ray emission from 3EG J1410–6147 may conceivably arise from a combination of PSRs J1412–6145, J1413–6141, and/or SNR G312.4–0.4.

The efficiency required to explain the EGRET flux in case #7 is  $\eta = 12\%$ , which seems possible. However, the spectral index of 2.50 is larger than those of known  $\gamma$ -ray pulsars. One of the SNRs coincident with the EGRET source, G337.8–0.1, harbors a maser (Koralesky et al. 1998), which is indicative of interaction between the SNR shock and the ambient medium. Thus, were a sufficiently massive molecular cloud located nearby, it could help produce the high energy radiation as a result of hadronic interaction (Aharonian, Drury, & Völk 1994; Aharonian & Atoyan 1996). Part of the EGRET flux could plausibly come from PSR J1637–4642 and part from pion  $\gamma$ -decay via SNR G337.8–0.1’s interaction with the putative cloud. In this case, the photon index would reflect a weighted average value. Both possible mechanisms for the high energy emission would produce a non-variable source, as is the case for 3EG J1639–4702.

Lastly we consider case #12, for which the required efficiency is high at the nominal pulsar distance and upper limit flux value,  $\eta = 55\%$ . The pulsar is located, in projection, just outside the plerionic SNR G27.8+0.6, for which the distance is  $\sim 2$  kpc (Reich et al. 1984). Although the estimated ages are comparable ( $\tau = 52$  kyr for the pulsar and  $\sim 45$  kyr for the SNR), it seems unlikely that both objects are physically associated, given the offset between the centrally peaked SNR component and the pulsar (see Reich et al. 1984). Whatever the possible relation between pulsar and SNR, the 3EG source in case #12 is variable (Table 2), arguing against a pulsar origin.

Examination of X-ray archives via HEASARC has revealed no compelling counterpart sources to any of the pulsars in Table 3. (A possible *Einstein* X-ray source at the edge of an IPC field near the location of PSR J1015–5719 [#3] is likely an artifact resulting from the large effective area correction used.) However, the exposures may not have been deep enough to reveal such X-ray emission, usually expected at the level  $L_x \sim 10^{-3}\dot{E}$  (e.g., Becker & Trumper 1997). Future observations of these

<sup>2</sup>To arrive at this result we have considered the 41 3EG sources in the range of the Parkes survey, and simulated them in synthetic populations in order to obtain an average expected number of chance coincidences. There is therefore a 98% a priori probability that not all 8 associations are spurious.

pulsar fields with *XMM-Newton* and *Chandra* could thus be instructive.

In conclusion, we find that two recently discovered Parkes pulsars (PSRs J1015–5719 and J1637–4642) could plausibly generate at least part of the  $\gamma$ -ray flux observed from two unidentified EGRET sources (3EG J1014–5705 and 3EG J1639–4702, respectively), and either of PSRs J1412–6145 or J1413–6141 could be associated with 3EG J1410–6147 if they are located closer than their DM distances by a factor of  $\sim 4$ . Cases #3–5 and 7 represent promising

targets for the forthcoming *AGILE* and *GLAST* missions.

D.F.T. was supported by CONICET and Fundación Antorchas, and is on leave from IAR, Argentina. Y.M.B. acknowledges the support of the High Energy Astrophysics division at the CfA and the *Chandra* project. Partial support from the US DOE grant DE-FG02-91ER40671 (D.F.T.) and NASA grants NAS8-39073 (Y.M.B.) and NAG5-9095 (F.C.) is acknowledged. We thank D. Nice and G. E. Romero for comments.

## REFERENCES

- Aharonian, F.A., Drury, L.O’C., & Völk, H.J. 1994, *A&A*, 285, 645  
 Aharonian, F.A., & Atoyan, A.M. 1996, *A&A*, 309, 917  
 Butt, Y., Torres, D.F., Combi, J.A., Dame, T., & Romero, G.E. 2001, in preparation  
 Camilo, F., et al. 2001, *ApJ*, 557, L51  
 Caraveo, P.A., De Luca, A., Mignani, R.P., Bignami, G.F. 2001 (*astro-ph/0107282*)  
 Case, G., & Bhattacharya, D., 1999, *ApJ* 521, 246  
 Caswell, J.L., & Barnes, P.J. 1985, *MNRAS*, 216, 753  
 Cheng, K.S., & Zhang, L., 1998, *ApJ*, 498, 327  
 Combi, J.A., Romero, G.E., & Azacárate, I. 1997, *Ap&SS*, 250, 1  
 Combi, J.A., & Romero, G.E. 1998a, *A&AS* 128, 423  
 Combi, J.A., Romero, G.E., & Benaglia P. 1998b, *A&A*, 333, L91  
 Combi, J.A., Romero, G.E., & Benaglia P. 1999, *ApJ*, 519, L177  
 Combi, J.A., Romero, G.E., Benaglia, P., & Jonas, J. 2001, *A&A*, 366, 1047  
 D’Amico, N., et al. 2001, *ApJ*, 552, L45  
 D’Amico, N. 1983, *Space Sci. Rev.*, 36, 195  
 Esposito, J.A., Hunter, S.D., Kanbach, G., & Sreekumar P. 1996, *ApJ*, 461, 820  
 Fierro, J.M., et al. 1993, *ApJ*, 413, L27  
 Green, D.A. 2000, *A Catalogue of Galactic Supernova Remnants*, Mullard Radio Astronomy Observatory, Cambridge, UK (available at <http://www.mrao.cam.ac.uk/surveys/snrs>)  
 Halpern, J.P., Camilo, F., Gotthelf, E.V., Helfand, D.J., Kramer, M., Lyne, A.G., Leighly, K.M., & Eracleous, M. 2001, *ApJ*, 552, L125  
 Hartman, R.C., et al. 1999, *ApJS*, 123, 79  
 Helfand, D.J. 1994, *MNRAS*, 267, 490  
 Kaspi, V.M., Lackey, J.R., Mattox, J., Manchester, R.N., Bailes, M., & Pace R. 2000, *ApJ*, 528, 445  
 Koralesky, B., Frail, D.A., Goss, W.M., Claussen, M.J., & Green, A.J. 1998, *AJ*, 116, 1323  
 Manchester, R.N., et al. 2001, *MNRAS*, in press (*astro-ph/0106522*)  
 Merck, M., et al., 1996, *A&AS*, 120, 365  
 Ögelman, H., & Finley, J.P. 1993, *ApJ*, 413, L31  
 Reich, W., Furst, E., Sofue, Y. 1984, *A&A*, 133, L4  
 Romero, G.E. 1998, *Rev. Mex. A&A*, 34, 29  
 Romero, G.E., Benaglia, P., & Torres D.F. 1999a *A&A*, 348, 868  
 Romero, G.E., Torres, D.F., Andruchow, I., Anchordoqui, L.A., & Link, B. 1999b, *MNRAS*, 308, 799  
 Ruiz, M.T., & May, J. 1986, *ApJ*, 309, 667  
 Sturmer, S.J., Dermer, C.D., & Mattox, J.R. 1996, *A&AS*, 120, 445  
 Taylor, J.H., & Cordes, J.M. 1993, *ApJ*, 411, 674  
 Thompson, D.J., 1999, in *International Symposium on High-Energy Gamma-Ray Astronomy*, Heidelberg, Germany, in press (*astro-ph/0101039*)  
 Thompson, D.J., et al. 1999, *ApJ*, 516, 297  
 Tompkins, W. 1999, Ph.D. Thesis, Stanford University  
 Torres, D.F., Romero, G.E., Combi, J., Benaglia, P., Punsly, B., & Andernach, H. 2001a, *A&A*, 370, 468  
 Torres, D.F., Combi, J.A., Romero, G.E., & Benaglia, P. 2001b, *Proc. Int. Workshop on The Nature of Galactic Unidentified Gamma-ray Sources*, O. Carramiñana, O. Reimer, D. Thomson (Eds.) (Kluwer Academic Press), 97  
 Torres, D.F., Pessah, M.E., & Romero, G.E. 2001c, *Astronomische Nachrichten* (2001), in press (*astro-ph/0104351*)  
 Yadigaroglu, I-A., & Romani, R.W. 1995, *ApJ*, 449, 211  
 Yadigaroglu, I-A., & Romani, R.W. 1997, *ApJ*, 476, 347  
 Zhang, L., Zhang, Y.J., & Cheng, K.S. 2000, *A&A*, 357, 957  
 Zhang, L., & Cheng, K.S. 1998, *A&A*, 335, 234

TABLE 1  
 $\gamma$ -RAY PULSARS DETECTED BY EGRET

| Pulsar/3EG Jsource | $P$<br>(ms) | $\tau$<br>(kyr) | $\dot{E}$<br>(ergs s $^{-1}$ ) | $d$<br>(kpc) | $F_{\gamma}^{3EG} [\times 10^{-8}]$<br>(ph cm $^{-2}$ s $^{-1}$ ) | $\gamma^{3EG}$  | $\eta$<br>(100MeV–<br>–10GeV) |
|--------------------|-------------|-----------------|--------------------------------|--------------|-------------------------------------------------------------------|-----------------|-------------------------------|
| Crab/0534–2200     | 33          | 1.2             | $5.0 \times 10^{38}$           | 2.0          | $226.2 \pm 4.7$                                                   | $2.19 \pm 0.02$ | 0.01%                         |
| Vela/0834–4511     | 89          | 12.5            | $6.3 \times 10^{36}$           | 0.25         | $834.3 \pm 11.2$                                                  | $1.69 \pm 0.01$ | 0.08%                         |
| B1951+32/ ...      | 39          | 100.0           | $3.7 \times 10^{36}$           | 2.4          | ...                                                               | ...             | 0.3%                          |
| B1706–44/1710–4439 | 102         | 15.8            | $3.1 \times 10^{36}$           | 1.8          | $111.2 \pm 6.2$                                                   | $1.86 \pm 0.04$ | 1%                            |
| Geminga/0633+1751  | 237         | 316.2           | $3.1 \times 10^{34}$           | 0.16         | $352.9 \pm 5.7$                                                   | $1.66 \pm 0.01$ | 3%                            |
| B1055–52/1058–5234 | 197         | 501.1           | $3.1 \times 10^{34}$           | 0.5/1.5      | $33.3 \pm 3.8$                                                    | $1.94 \pm 0.10$ | 2/19%                         |

NOTE.—Pulsar parameters and distances (and  $\eta$  in the case of PSR B1951+32 — this pulsar is not a 3EG source) are taken from Kaspi et al. (2000), excepting PSR B1055–52, for which we also consider a smaller value of distance (Ögelman & Finley 1993; Combi et al. 1997), and Vela (Caraveo et al. 2001 and references therein).  $\tau = P/2\dot{P}$ , and  $\dot{E} = 4\pi^2 I \dot{P}/P^3$ , with  $I = 10^{45}$  g cm $^2$ . The “P1234”  $\gamma$ -ray fluxes and spectral indices are from the 3EG catalog (Hartman et al. 1999), from which we have computed the  $\eta$  values.

TABLE 2  
POSITIONAL COINCIDENCES BETWEEN UNIDENTIFIED 3EG SOURCES AND PARKES PULSARS

| 3EG J     | Note  | $I$ | PSR J     | #  | SNRs                               | Ref.                  | $\dot{E}/d^2$        |
|-----------|-------|-----|-----------|----|------------------------------------|-----------------------|----------------------|
| 1013–5915 | C, em | 1.6 | 1016–5857 | 1  | G284.3–1.8                         | Camilo et al. (2001)  | $2.9 \times 10^{35}$ |
|           |       |     | 1013–5934 | 2  |                                    |                       | $2.0 \times 10^{30}$ |
| 1014–5705 | C, em | 1.4 | 1015–5719 | 3  |                                    |                       | $3.4 \times 10^{34}$ |
| 1410–6147 | C     | 1.2 | 1412–6145 | 4  | G312.4–0.4                         |                       | $1.4 \times 10^{33}$ |
|           |       |     | 1413–6141 | 5  |                                    |                       | $1.6 \times 10^{33}$ |
| 1420–6038 | C     | 2.1 | 1420–6048 | 6  |                                    | D’Amico et al. (2001) | $1.7 \times 10^{35}$ |
| 1639–4702 | C, em | 1.9 | 1637–4642 | 7  | G337.8–0.1; G338.1+0.4; G338.3+0.0 |                       | $1.9 \times 10^{34}$ |
|           |       |     | 1640–4648 | 8  |                                    |                       | $1.5 \times 10^{32}$ |
|           |       |     | 1637–4721 | 9  |                                    |                       | $3.1 \times 10^{30}$ |
| 1714–3857 | C, em | 2.1 | 1713–3844 | 10 | G348.5+0.0; G348.5+0.1; G347.3–0.5 | Butt et al. (2001)    | $4.0 \times 10^{31}$ |
|           |       |     | 1715–3903 | 11 |                                    | Butt et al. (2001)    | $3.0 \times 10^{33}$ |
| 1837–0423 | C     | 5.4 | 1838–0453 | 12 | G27.8+0.6                          |                       | $1.2 \times 10^{33}$ |
| 1837–0606 | C, em | 2.4 | 1837–0559 | 13 |                                    |                       | $6.5 \times 10^{32}$ |
|           |       |     | 1837–0604 | 14 |                                    | D’Amico et al. (2001) | $5.2 \times 10^{34}$ |

NOTE.—A number (#) is assigned to each pair for ease of reference. SNRs contained in Green’s (2000) catalog coinciding with the *EGRET* sources are noted (see Torres et al. 2001b). Only in cases #4, 5 and 12 do the pulsars also coincide with the SNRs listed. “C” and “em” refer to the  $\gamma$ -ray sources: source confusion exists and sources are possibly extended or multiple, respectively (Hartman et al. 1999).  $I$  is the variability index as in Torres et al. (2001a, c), where  $I > 5$  ( $< 2$ ) represents a source whose flux presents variability levels at least  $8\sigma$  (less than  $2\sigma$ ) above those displayed by confirmed pulsars. Pulsar “spin-down flux”  $\dot{E}/d^2$  is in units of  $\text{ergs s}^{-1} \text{ kpc}^{-2}$ .

TABLE 3  
OBSERVATIONAL PARAMETERS FOR THE PULSARS IN THE CANDIDATE ASSOCIATIONS

| Case | $\Delta\theta$ [deg] | $\theta$ [deg] | $(l, b)$      | $d$ [kpc] | $\tau$ [kyr] | $P$ [ms] | $\dot{P}$ [ $10^{-15}$ ] | $\dot{E}$ [ $\text{ergs s}^{-1}$ ] |
|------|----------------------|----------------|---------------|-----------|--------------|----------|--------------------------|------------------------------------|
| #3   | 0.30                 | 0.67           | 283.09, –0.58 | 4.9       | 38.7         | 140      | 57.4                     | $8.2 \times 10^{35}$               |
| #4   | 0.15                 | 0.36           | 312.32, –0.37 | 9.3       | 50.6         | 315      | 98.7                     | $1.2 \times 10^{35}$               |
| #5   | 0.28                 | 0.36           | 312.46, –0.34 | 11.0      | 13.5         | 286      | 333.4                    | $5.7 \times 10^{35}$               |
| #7   | 0.46                 | 0.56           | 337.79, +0.31 | 5.8       | 41.2         | 154      | 59.2                     | $6.4 \times 10^{35}$               |
| #12  | 0.50                 | 0.52           | 27.07, +0.71  | 8.2       | 52.2         | 381      | 115.7                    | $8.3 \times 10^{34}$               |

NOTE.— $\Delta\theta$  is the angular distance between the center of the 3EG source and the position of the respective coincident pulsar.  $\theta$  is the effective 95% confidence level radius of the 3EG source error box (Hartman et al. 1999). Pulsar parameters are taken from the Parkes survey database (see footnote 1), and their distances ( $d$ ) are estimated from the observed dispersion measure (Taylor & Cordes 1993).

TABLE 4  
OBSERVED AND COMPUTED  $\gamma$ -RAY EMISSION PARAMETERS FOR NEW CANDIDATE ASSOCIATIONS

| Case | $F^{3EG} [\times 10^{-8} \text{ ph cm}^{-2} \text{ s}^{-1}]$ | $F^{3EG} [\text{ergs cm}^{-2} \text{ s}^{-1}]$ | $\gamma^{3EG}$  | $L_{isotropic} [\text{ergs s}^{-1}]$ | $L_{beamed} [\text{ergs s}^{-1}]$ | $\eta$               |
|------|--------------------------------------------------------------|------------------------------------------------|-----------------|--------------------------------------|-----------------------------------|----------------------|
| #3   | $34.0 \pm 6.5$                                               | $1.90 \times 10^{-10}$                         | $2.23 \pm 0.20$ | $5.2 \times 10^{35}$                 | $4.1 \times 10^{34}$              | 5%                   |
| #4   | $64.2 \pm 8.8$                                               | $4.09 \times 10^{-10}$                         | $2.12 \pm 0.14$ | $4.1 \times 10^{36}$                 | $3.2 \times 10^{35}$              | 270/12% <sup>a</sup> |
| #5   | $64.2 \pm 8.8$                                               | $4.09 \times 10^{-10}$                         | $2.12 \pm 0.14$ | $5.7 \times 10^{36}$                 | $4.5 \times 10^{35}$              | 80/3% <sup>a</sup>   |
| #7   | $53.2 \pm 8.7$                                               | $2.30 \times 10^{-10}$                         | $2.50 \pm 0.18$ | $9.3 \times 10^{35}$                 | $7.4 \times 10^{34}$              | 12%                  |
| #12  | $< 19.1$                                                     | $< 0.70 \times 10^{-10}$                       | $2.71 \pm 0.44$ | $< 5.8 \times 10^{35}$               | $< 4.6 \times 10^{34}$            | $< 55\%$             |

NOTE.—The EGRET fluxes correspond to the “P1234” values (Hartman et al. 1999). Luminosities, and efficiency values for all pulsars, are given for the distance estimated from the dispersion measure (Table 3).

<sup>a</sup>Second values of efficiency correspond to a distance of 1.9 kpc, one of those estimated for the positionally coincident SNR G312.4–0.4 (see text).